

Colonoscopy with SUPREP SPLIT DOSE

		• •	
Pati	ient Name:	Physician Name:	Today's Date:
Date of Procedure: Proce		Procedure Time:	Arrival Time:
Pro	cedure Location:		
GENERAL INFORMATION:			5 – 7 DAYS BEFORE YOUR PROCEDURE:
>	IV sedation or anesthesia is usually given during your procedure. Due to regulatory requirements and for your safety, you must arrange for an adult (18+) to drive you home, or risk procedure cancellation.		IF YOU HAVE DIABETES: Follow your GI Physician's recommendations for oral and insulin medication.
>	> You will not be able to drive, operate machinery, make important decisions, or return to work for the rest of the day.		If you are on Aspirin, Coumadin, Warfarin, Plavix, Heparin, Lovenox, Fish Oil, or any otherblood thinners:

- You may resume normal activities the next day unless the doctor states otherwise.
- > If you have any questions concerning your procedure or need to cancel, please call the respective office.
- > Wear comfortable, loose-fitting clothing that is easy to step into. Wear flat shoes or tennis shoes.
- > The center is not responsible for your property. Do not wear jewelry orbring valuables with you to the Center or Hospital.
- The morning of the procedure, only drink the required liquid preparation.

YOU WILL NEED TO PURCHASE: Clenpiq – Prescription Medication

- Your gastroenterologist may have advisedyou to stop taking blood thinners prior to your procedure. You will need to stop taking these medications 5-7 days prior to your procedure.
- The GI physician may have to obtain Cardiac Clearance before your procedurecan be scheduled. The office will contact you once clearance is obtained.
- > 5-7 days before the procedure, stop eating salads, raw vegetables, nuts, seeds, and popcorn.



WHAT YOU CANNOT DRINK



- Any liquid that is red, orange, blue, or purple in color
- No milk, dairy cream, or non-dairy products
- No alcohol
- No orange juice or lemonade

WHAT YOU CAN DRINK

- White grape juice, white cranberry juice, and apple juice
- > Kool-Aide, Powerade, Gatorade, and Crystal Light
- Popsicles and Italian Ice
- Broth or bouillon: beef, chicken, or vegetable flavored
- Water, soda, tea, or coffee
- Hard candy and sugar
- Jell-O

DO NOT consume red, orange, blue or purple liquid from this list

THE KEY TO A GOOD PREPERATION

Drink plenty of the correct fluids, see 'What You Can Drink' list above. You must take <u>ALL</u> of the PREP solution as directed to clean your bowel adequately. Even if you are passing clear liquid, you must continue to take the entire dose of the prep the day before your exam. Failure to follow prep instructions may result in rescheduling of the procedure due to poor preparation.

THE DAY BEFORE YOUR COLONOSCOPY

- No solid food and NO ALCOHOL
- Clear liquids only all day
- 8 oz. = 1 Cup and 16 oz. = 2 Cup

<u>5 PM – 6 PM:</u> Pour the entire contents of (1 of 2) bottles of SUPREP Bowel Prep Kit into the mixing container provided. Fill the container with water to the 16oz. **fill line** as indicated on the mixing container and drink the entire amount.

- Continue to drink 16 to 32 oz. of clear liquids until you go to bed. Please refer to the 'What You Can Drink' list in the first column. No red, orange, blue, or purple liquid.
- Apply a petroleum-based product or diaper rash ointment to the rectal area if you experience discomfort from frequent stools.
- Confirm that you have a driver to take you home from your procedure.



THE DAY OF YOUR COLONOSCOPY

You should take your essential medications with a small sip of water (e.g. heart and blood pressure).

7 hours prior to procedure, pour the entire contents of (2 of 2) bottles of SUPREP Bowel Prep Kit into the mixing container provided. Fill the container withwater to the 16oz. **fill line** as indicated on the mixing container and drink the entire amount.

- Following consumption of the prep,drink an additional 32 oz. of water within an hour.
- No liquids 4 hours before procedure.



Additional Comments: